

N 832

Seat No.

--	--	--	--	--	--

2025 III 07 1100 – N 832– MATHEMATICS (71) GEOMETRY—PART II (E)

(REVISED COURSE)

Time : 2 Hours

(Pages 12)

Max. Marks : 40

Note :—

- (i) All questions are compulsory.
- (ii) Use of a calculator is not allowed.
- (iii) The numbers to the right of the questions indicate full marks.
- (iv) In case of MCQs [Q. No. 1(A)] only the first attempt will be evaluated and will be given credit.
- (v) Draw proper figures wherever necessary.
- (vi) The marks of construction should be clear. Do not erase them.
- (vii) Diagram is essential for writing the proof of the theorem.

1. (A) Choose the correct alternative from given : 4

(1) Out of the following which is a Pythagorean triplet ?

(A) (1, 5, 10)

(B) (3, 4, 5)

(C) (2, 2, 2)

(D) (5, 5, 2)

P.T.O.

2/N 832

- (2) $\angle ACB$ is inscribed angle in a circle with centre O. If $\angle ACB = 65^\circ$, then what is measure of its intercepted arc AXB ?

(A) 65°

(B) 230°

(C) 295°

(D) 130°

- (3) Distance of point (3, 4) from the origin is

(A) 7

(B) 1

(C) 5

(D) -5

- (4) If radius of cone is 5 cm and its perpendicular height is 12 cm, then the slant height is

(A) 17 cm

(B) 4 cm

(C) 13 cm


(D) 60 cm

3/N 832


(B) Solve the following sub-questions : 4

(1) In the following figure $\triangle ABC$, $B - D - C$ and $BD = 7$, $BC = 20$,

then find $\frac{A(\triangle ABD)}{A(\triangle ABC)}$.



(2) In the following figure $\angle MNP = 90^\circ$, seg $NQ \perp$ seg MP , $MQ = 9$, $QP = 4$, find NQ .



(3) Angle made by a line with the positive direction of X-axis is 30° . Find slope of that line.

(4) In cyclic quadrilateral ABCD $m\angle A = 100^\circ$, then find $m\angle C$.


P.T.O.

4/N 832

2. (A) Complete the following activities and rewrite it (any *two*) : 4

- (1) The radius of a circle with centre 'P' is 10 cm. If chord AB of the circle subtends a right angle at P, find area of minor sector

by using the following activity. ($\pi = 3.14$)



Activity :

$$r = 10 \text{ cm}, \theta = 90^\circ, \pi = 3.14.$$


$$\begin{aligned} A(P\text{-}AXB) &= \frac{\theta}{360} \times \boxed{} \\ &= \frac{\boxed{}}{360} \times 3.14 \times 10^2 \end{aligned}$$

$$= \frac{1}{4} \times \boxed{}$$

$$A(P\text{-}AXB) = \boxed{} \text{ sq. cm.}$$

5/N 832

- (2) In the following figure chord MN and chord RS intersect at point D. If RD = 15, DS = 4, MD = 8, find DN by completing the following activity :



Activity :

$$\therefore MD \times DN = \boxed{} \times DS \dots\dots\dots$$

..... (Theorem of internal division of chords)

$$\therefore \boxed{} \times DN = 15 \times 4$$


$$\therefore DN = \frac{\boxed{}}{8}$$

$$\therefore DN = \boxed{}$$

P.T.O.

6/N 832

- (3) An observer at a distance of 10 m from tree looks at the top of the tree, the angle of elevation is 60° . To find the height of tree complete the activity. ($\sqrt{3} = 1.73$)



Activity :

In the figure given above, $AB = h$ = height of tree, $BC = 10$ m, distance of the observer from the tree.

Angle of elevation (θ) = $\angle BCA = 60^\circ$

$$\tan \theta = \frac{\boxed{}}{BC} \dots\dots\dots \text{(I)}$$

$$\tan 60^\circ = \boxed{} \dots\dots\dots \text{(II)}$$

$$\frac{AB}{BC} = \sqrt{3} \quad (\text{From (I) and (II)})$$

$$AB = BC \times \sqrt{3} = 10\sqrt{3}$$

$$AB = 10 \times 1.73 = \boxed{}$$


$\therefore$ height of the tree is $\boxed{}$ m.

7/N 832


(B) Solve the following sub-questions (any *four*) :

8

- (1) In $\triangle ABC$, $DE \parallel BC$. If $DB = 5.4$ cm, $AD = 1.8$ cm, $EC = 7.2$ cm, then find AE .




- (2) In the figure given below, find RS and PS using the information given in $\triangle PSR$.



P.T.O.

8/N 832

- (3) In the following figure, circle with centre D touches the sides of $\angle ACB$ at A and B. If $\angle ACB = 52^\circ$, find measure of $\angle ADB$.




- (4) Verify, whether points, A(1, -3), B(2, -5) and C(-4, 7) are collinear or not.

- (5) If $\sin \theta = \frac{11}{61}$, find the values of $\cos \theta$ using trigonometric identity.

3. (A) Complete the following activities and rewrite it (any one) : 3

- (1) In the following figure, $XY \parallel \text{seg } AC$. If $2AX = 3BX$ and $XY = 9$. Complete the activity to find the value of AC.



9/N 832

Activity :

$$2AX = 3BX \text{ (Given)}$$

$$\therefore \frac{AX}{BX} = \frac{3}{\boxed{}}$$

$$\frac{AX + BX}{BX} = \frac{3 + 2}{2} \text{ (by componendo)}$$

$$\frac{\boxed{}}{BX} = \frac{5}{2} \text{ (I)}$$


Now $\triangle BCA \sim \triangle BYX$ ($\boxed{}$ test of similarity)

$$\therefore \frac{BA}{BX} = \frac{AC}{XY} \text{ (corresponding sides of similar triangles)}$$

$$\frac{\boxed{}}{\boxed{}} = \frac{AC}{9} \text{ from (I)}$$

$$\therefore AC = \boxed{}$$

- (2) Complete the following activity to prove that the sum of squares of diagonals of a rhombus is equal to the sum of the squares of the sides.



P.T.O.

10/N 832

Given :

□ PQRS is a rhombus. Diagonals PR and SQ intersect each other at point T.

To prove : $PS^2 + SR^2 + QR^2 + PQ^2 = PR^2 + QS^2$

Activity :

Diagonals of a rhombus bisect each other.

In ΔPQS , PT is the median and in ΔQRS , RT is the median.

$\therefore$ by Apollonius theorem,

$$PQ^2 + PS^2 = \boxed{} + 2QT^2 \dots\dots\dots (I)$$

$$QR^2 + SR^2 = \boxed{} + 2QT^2 \dots\dots\dots (II)$$

adding (I) and (II),

$$PQ^2 + PS^2 + QR^2 + SR^2 = 2(PT^2 + \boxed{}) + 4QT^2$$

$$= 2(PT^2 + \boxed{}) + 4QT^2$$

$$\dots\dots\dots (RT = PT)$$

$$= 4PT^2 + 4QT^2$$

$$= (\boxed{})^2 + (2QT)^2$$

$$\therefore PQ^2 + PS^2 + QR^2 + SR^2 = PR^2 + \boxed{}.$$

11/N 832

(B) Solve the following sub-questions (any *two*) :

6


- (1) Show that points $P(1, -2)$, $Q(5, 2)$, $R(3, -1)$, $S(-1, -5)$ are the vertices of a parallelogram.
- (2) Prove that tangent segments drawn from an external point to a circle are congruent.
- (3) Draw a circle with radius 4.1 cm. Construct tangents to the circle from a point at a distance 7.3 cm from the centre.
- (4) How many solid cylinders of radius 10 cm and height 6 cm can be made by melting a solid sphere of radius 30 cm ?

4. Solve the following sub-questions (any *two*) :

8

(1) In the following figure $DE \parallel BC$, then :

- (i) If $DE = 4$ cm, $BC = 8$ cm, $A(\triangle ADE) = 25$ cm², find $A(\triangle ABC)$.
- (ii) If $DE : BC = 3 : 5$, then find $A(\triangle ADE) : A(\square DBCE)$.



P.T.O.

12/N 832

- (2) $\Delta ABC \sim \Delta PQR$. In ΔABC , $AB = 3.6$ cm, $BC = 4$ cm and $AC = 4.2$ cm.

The corresponding sides of ΔABC and ΔPQR are in the ratio $2 : 3$,
construct ΔABC and ΔPQR .

- (3) The radii of the circular ends of a frustum of a cone are 14 cm and 8 cm. If the height of the frustum is 8 cm, find : ($\pi = 3.14$)

- (i) Curved surface area of frustum.
- (ii) Total surface area of the frustum.
- (iii) Volume of the frustum.

5. Solve the following sub-questions (any one) :

3

- (1) $\square ABCD$ is a rectangle. Taking AD as a diameter, a semicircle AXD is drawn which intersects the diagonal BD at X . If $AB = 12$ cm, $AD = 9$ cm, then find the values of BD and BX .

- (2) Taking $\theta = 30^\circ$ to verify the following Trigonometric identities :

- (i) $\sin^2 \theta + \cos^2 \theta = 1$
- (ii) $1 + \tan^2 \theta = \sec^2 \theta$
- (iii) $1 + \cot^2 \theta = \operatorname{cosec}^2 \theta$.